









Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

Good Poultry Management

Rules for Every-Day Use:

1. Buy good stock.
2. Keep buildings and equipment clean.
3. Provide proper heat and ventilation.
4. Supply plenty of Southern States feed – clean water always.

KIND OF POULTRY	FLOOR SPACE	WATER SPACE	FEEDING SPACE	ROOSTING SPACE
 BABY CHICKS	1/2 square ft. to 6 weeks	1 1-gallon fount to each 50	1 inch apiece first 3 weeks 2 inches each, 3 to 6 weeks	Supply at 3 weeks
 GROWING PULLETS	1-3 square ft. each	1 3-gallon fount to each 50	3 inches each; plenty of room for grain if fed	5-8 running inches apiece
 COCKERELS	Same as Pullets	Same as Pullets	Same as Pullets	Not needed if being fattened
 ** LAYERS AND BREEDERS	4 square ft. apiece	1 5-gallon fount to each 50	4 running in. each. 20-25 birds to one hanging hopper	6-9 inches apiece
 BROILERS	1 square ft. apiece	Same as Pullets	Same as Pullets	Not needed
 TURKEY POULTS	1 square ft. to 6 weeks	2 1-gallon founts to each 50	2 inches each, first 3 weeks	Supply at 3 weeks
 GROWING TURKEYS	3-5 square ft. apiece	2 5-gallon founts to each 50	3-6 inches apiece	12-15 inches apiece
 † TURKEYS LAYERS AND BREEDERS	8 square ft. apiece	2 5-gallon founts to each 50	8 inches each	15 inches apiece

* * 15 nests for each 100 birds—size: 15x15

† 20 large nests (20x20) for each 100 birds.

SOUTHERN STATES COOPERATIVE



General Offices:
RICHMOND, VA.